



One hospital, one appendectomy: The cost effectiveness of a standardized doctor's preference card[☆]



David E. Skarda^{a,*}, Michael Rollins^a, Seth Andrews^b, Molly McFadden^c, Doug Barnhart^a, Rebecca Meyers^a, Eric Scaife^a

^a Division of Pediatric Surgery, University of Utah, Primary Children's Hospital, Salt Lake City, UT

^b Systems Improvement, Primary Children's Hospital, Salt Lake City, UT

^c Division of Epidemiology, University of Utah School of Medicine, Salt Lake City, UT

ARTICLE INFO

Article history:

Received 28 February 2015

Accepted 10 March 2015

Key words:

Appendicitis

Value

Pediatric

Laparoscopy

Cost

Standardize

ABSTRACT

Purpose: Appendicitis in children provides a unique opportunity to explore changes that reduce variation, reduce cost, and improve value. In this study we sought to evaluate the effectiveness of standardization of surgical technique and intraoperative disposable device utilization for laparoscopic appendectomy among all surgeons at a tertiary children's hospital.

Methods: All 6 surgeons at our tertiary children's hospital agreed to standardize to a single technique of performing a laparoscopic appendectomy. We collected data on all pediatric patients who had a laparoscopic appendectomy following implementation of the uniform doctor's preference card (DPC) (March 1, 2013 to February 28, 2014) and compared them to a historical control group.

Results: Implementation of the uniform DPC decreased the device cost per appendectomy from \$844.11 to \$305.32. Operative times (skin incision to skin closure) were 34.8 minutes prior to the uniform DPC and 37.0 minutes using the uniform DPC. There were no significant differences in postappendectomy outcomes.

Conclusion: We have demonstrated that implementation of a uniform DPC and technical standardization for laparoscopic appendectomy can significantly reduce cost. Furthermore, this can occur without dramatically increasing operative times, length of stay, or postoperative complications.

© 2015 Elsevier Inc. All rights reserved.

Variation in laparoscopic appendectomy technique, as with many surgical procedures, is common and is based on individual surgeon training and preference. Despite differences in surgical technique among surgeons, outcomes are relatively similar. Given the high clinical volume combined with the low frequency of complications, appendicitis represents a unique opportunity to increase value by implementing cost-saving measures, reducing variability and minimizing resource utilization. In this study we sought to evaluate the effectiveness of standardization of surgical technique and intraoperative disposable device utilization for laparoscopic appendectomy among 6 surgeons at a tertiary children's hospital.

1. Methods

After obtaining institutional review board (IRB) (IRB no. 00064063) approval we performed a retrospective query of a prospectively maintained database at our tertiary care children's hospital. We standardized

the doctor's preference card (DPC) for laparoscopic appendectomy on March 1, 2013.

1.1. Prior to uniform DPC

Prior to the implementation of the uniform DPC, 6 board certified pediatric surgeons at our children's hospital performed laparoscopic appendectomies using various techniques. Instruments varied from surgeon to surgeon but often included dilating sheathed ports, linear stapler(s), a pistol-grip heat source and a specimen retrieval bag.

1.2. Uniform DPC

We created and implemented the uniform DPC for laparoscopic appendectomy by consensus opinion among the 6 surgeons and based the selection of devices in our uniform DPC on cost, availability and utility. We identified several effective and inexpensive devices available in our healthcare system but not previously utilized at our institution. Changes regarding the instruments and technique included low-cost dilating ribbed ports, use of a reusable hook cautery to divide the mesoappendix and pre-tied surgical loops to control the base of the appendix and cecum. The uniform DPC excluded linear staplers, pistol-grip heat sources and specimen retrieval bags. These were only used if the

[☆] Support: None.

* Corresponding author at: Primary Children's Hospital, 100 North Mario Capecchi Drive, Suite 2600, Salt Lake City, UT, 84113. Tel.: +1 801 662 2950; fax: +1 801 662 2980.

E-mail address: david.skarda@gmail.com (D.E. Skarda).

operating surgeon determined during the procedure that the device was necessary.

1.3. Definition of ruptured vs. nonruptured appendicitis

We strictly defined ruptured appendicitis as a visible hole in the appendix or a fecalith in the abdomen [1]. Patients with gangrenous appendicitis were included in this analysis as nonruptured appendicitis in both analysis groups. The nonruptured status was based upon intraoperative identification and did not change during the postoperative hospital course. Pathologic results did not affect how patients were categorized or treated postoperatively. If the appendix was ruptured intraoperatively owing to surgical manipulation, that patient was categorized as “ruptured”.

1.4. Activity-based accounting

Costs were derived from the hospital's cost-accounting program, the Standard Cost Master, which is a transaction-based microcosting system. This system identifies and aggregates the variable- and fixed-cost components of patient activities, hospital services, and products according to the date of service [2,3]. Total hospital costs are shown. Hospital costs incurred in years 2011–2014 were standardized to 2014 US dollars by applying a yearly consumer price index for hospital services [4]. Cost reported in this analysis includes the hospital cost of the initial evaluation, procedure, and hospital stay.

1.5. Statistics

Univariate analyses comparing before to after uniform DPC procedures and outcomes were performed. For categorical variables, differences in proportions between the 2 time periods were tested with the chi square test or Fisher's Exact test as appropriate. Continuous variable differences were tested using ANOVA or the equivalent nonparametric procedure. These analyses were performed using SAS version 9.4 (2013; SAS Institute Inc., Cary, NC).

Costs were converted to 2014 US dollars using the Consumer Price Index. Differences in the costs across the two study periods were tested using a multivariable generalized linear model (GLM) with a log link function. While the gamma distribution is typically used in GLM regressions for healthcare cost outcomes, the GLM family test (modified Park test) indicated that the most appropriate distribution for these data was the Poisson. Results were returned to the original cost scale using the margins command in Stata. These analyses were performed using Stata version 13 (2013; StataCorp LP, College Station, TX).

Significance was defined as P value $\leq .05$. Descriptive statistics were calculated as mean with 95% confidence interval or N (%).

2. Results

From March 1, 2013 to February 28, 2014 362 consecutive laparoscopic appendectomies were performed using the uniform DPC. In the historical control group there were 346 consecutive laparoscopic appendectomies performed prior to the implementation of the uniform DPC from March 1, 2012 to February 28, 2013. Comparisons of patients with ruptured and nonruptured appendicitis in whom the appendectomies were performed using the uniform DPC vs. the historical control group reveal no differences in gender, age, or WBC count on admission (Table 1).

2.1. Effect of the uniform DPC on the cost of the procedure

Implementation of the uniform DPC decreased the device cost per appendectomy from \$829.73 to \$279.76 for patients with nonruptured appendicitis and from \$874.08 to \$361.51 for patients with ruptured appendicitis. The cost savings directly attributable to the implementation of the uniform DPC during the one year time period of this study was \$195,041.98. Operative times (skin incision to skin closure) were

Table 1
Demographic, cost, and duration information.

	Prior to uniform DPC, N = 346	With uniform DPC, N = 362	P value
	Mean (95% CI)	Mean (95% CI)	
All patients with appendicitis			
Age in years	9.4 (9.0–9.8)	10.0 (9.6–10.4)	0.047
Gender (M:F)	188:158 (54.3% Male)	196:166 (54.1% Male)	0.959
Admission WBC	13.8 (13.2–14.5)	13.7 (13.1–14.4)	0.758
Appendectomy device cost	\$844.11 (838.65–849.56)	\$305.32 (302.05–308.60)	<0.001
Skin to skin	34.8 (33.3–36.3)	37.0 (35.6–38.4)	0.006
OR time (min)			
Roll-in roll-out	59.3 (57.6–61.1)	61.7 (60.0–63.3)	0.016
OR time (min)			
Length of stay (hours)	50.9 (44.1–57.7)	46.7 (41.1–52.3)	0.394
Patients with nonruptured appendicitis			
	N = 229	N = 247	
Age in years	10.0 (9.6–10.5)	10.5 (10.1–10.9)	0.047
Gender (M:F)	123:106 (53.7% Male)	139:108 (56.3% Male)	0.574
Admission WBC	13.4 (12.6–14.2)	14.2 (13.3–15.0)	0.139
Appendectomy device cost	\$829.73 (823.63–835.84)	\$279.76 (276.22–283.29)	<0.001
Skin to skin OR time (min)	31.2 (30.2–33.4)	34.0 (32.3–35.3)	0.006
Roll-in roll-out OR time (min)	55.6 (53.7–57.5)	58.3 (56.6–60.0)	0.016
Length of stay (hours)	22.5 (20.3–24.6)	24.9 (20.9–28.9)	0.394
Patients with ruptured appendicitis			
	N = 117	N = 115	
Age in years	8.2 (7.5–8.9)	8.9 (8.2–9.6)	0.047
Gender (M:F)	65:52 (55.6% Male)	57:58 (49.6% Male)	0.361
Admission WBC	14.4 (13.3–15.5)	13.0 (11.9–14.0)	0.055
Appendectomy device cost	\$874.08 (867.69–880.48)	\$361.51 (357.41–365.61)	<0.001
Skin to skin OR time (min)	41.7 (38.9–44.4)	43.4 (40.6–46.3)	0.006
Roll-in roll-out OR time (min)	66.6 (63.4–69.9)	68.8 (65.3–72.4)	0.016
Length of stay (hours)	106.6 (91.2–121.7)	93.6 (82.2–104.9)	0.394

31.2 minutes prior to the uniform DPC and 34.0 minutes using the uniform DPC for patients with nonruptured appendicitis and 41.7 minutes versus 43.4 minutes for patients with ruptured appendicitis. Total time in the operating room was 55.6 minutes prior to the uniform DPC and 58.3 minutes using the uniform DPC for patients with nonruptured appendicitis and 66.6 minutes versus 68.8 minutes for patients with ruptured appendicitis (Table 1).

2.2. Effect of the uniform DPC on the overall cost of the treatment of appendicitis

Hospital length of stay increased from 22.5 hours prior to the uniform DPC to 24.9 hours using the uniform DPC in patients with nonruptured appendicitis and decreased from 106.6 hours prior to the uniform DPC to 93.6 hours using the uniform DPC in patients with ruptured appendicitis (Table 1).

2.3. Effect of the uniform DPC on outcomes

There were no significant differences in postappendectomy outcomes including inpatient readmission, observation readmission, postdischarge emergency department reevaluation, abdominal abscess, reoperation, postoperative interventional radiology drainage, subcutaneous abscess,

Table 2
Outcomes.

	Prior to uniform DPC N = 346	With uniform DPC N = 362		P value
	12 month period	12 month period	First 6 months	
	N (%)	N (%)	N	
All patients with appendicitis				
Inpatient readmission	8 (2.3%)	14 (%)	11	0.250
Observation readmission	16 (4.6%)	14 (%)	6	0.759
Postdischarge emergency department evaluation	12 (3.5%)	13 (%)	7	0.179
Abdominal abscess	4 (1.2%)	9 (2.5%)	8	0.413
Reoperation	1 (0.3%)	3 (0.8%)	2	0.624
Interventional radiology drainage	3 (0.9%)	6 (1.7%)	6	0.509
Subcutaneous abscess	3 (0.9%)	4 (1.1%)	3	1.000
<i>C. difficile</i> colitis	1 (0.3%)	3 (0.8%)	1	0.624
Postoperative CT	23 (6.6%)	19 (5.2%)	14	0.431
Patients with nonruptured appendicitis				
Inpatient readmission	3 (1.3%)	5 (1.6%)	3	1.000
Observation readmission	6 (2.6%)	8 (3.2%)	3	1.000
Postdischarge emergency department evaluation	8 (3.5%)	8 (4.0%)	2	0.155
Abdominal abscess	1 (0.4%)	3 (1.2%)	2	0.413
Reoperation	0 (0.0%)	2 (0.8%)	1	0.500
Interventional radiology drainage	1 (0.4%)	1 (0.4%)	1	1.000
Subcutaneous abscess	1 (0.4%)	1 (0.4%)	1	1.000
<i>C. difficile</i> colitis	0 (0.0%)	3 (1.2%)	1	0.249
Postoperative CT	12 (5.2%)	11 (4.5%)	7	0.689
Patients with ruptured appendicitis				
Inpatient readmission	5 (4.3%)	9 (%)	8	0.223
Observation readmission	10 (8.5%)	6 (%)	4	0.729
Postdischarge emergency department evaluation	4 (3.4%)	5 (%)	4	0.747
Abdominal abscess	3 (2.6%)	6 (5.2%)	6	0.278
Reoperation	1 (0.9%)	1 (0.9%)	1	1.000
Interventional radiology drainage	2 (1.7%)	5 (4.3%)	5	0.278
Subcutaneous abscess	2 (1.7%)	3 (2.6%)	2	0.682
<i>C. difficile</i> colitis	1 (0.9%)	0 (0.0%)	0	1.000
Postoperative CT	11 (9.4%)	8 (7.0%)	7	0.497

Clostridium difficile colitis, and postoperative CT imaging between patients treated before and after implementation of the uniform DPC. Although there were generally more complications after implementation of the uniform DPC, most of these events occurred during the first 6 months after implementation (Table 2).


2.4. Effect of the uniform DPC on between surgeon variability

Surgeon variability of appendectomy procedure device cost decreased with implementation of the uniform DPC (Figs. 1 and 2).

Uniform DPC

No. Device

- 1 chlorhexidine-based skin prep
- 1 sterile drapes
- 4 sterile gowns
- 4 sterile gloves
- 1 local anesthetic
- 1 standard laparoscopy set with tower and insufflation tubing
- 2 5 mm port
- 1 12 mm port
- 2 0 pre-tied loops
- 1 2-0 fascia closure suture
- 1 5-0 skin closure suture
- 3 surgical site dressing

Fig. 1. Uniform DPC.**Fig. 2.** Between surgeon variability is surgical device cost per appendectomy.

2.5. Effect of the uniform DPC on specific device utilization

Sheathed port utilization decreased from use in 99.2% of laparoscopic appendectomies prior to the uniform DPC to 9.6% of laparoscopic appendectomies using the uniform DPC. Nonsheathed low-cost port utilization increased from use in 0.0% prior to the uniform DPC to use in 92.6% of laparoscopic appendectomies with the uniform DPC. Stapler utilization decreased from use in 94.4% of laparoscopic appendectomies to 20.3% using the uniform DPC. Low cost pre-tied loop utilization increased from use in 0.0%–79.7% of laparoscopic appendectomies with the uniform DPC. Pistol-grip heat source utilization decreased from use in 22.3%–3.8% using the uniform DPC (Table 3).

3. Discussion

Appendicitis remains the most common indication for urgent surgical intervention in children. Optimizing resource utilization for these patients has the potential to significantly reduce cost and improve value. Clinical protocols have been shown to reduce costs in pediatric appendicitis [5]. A single institution study demonstrated a 20% reduction in cost with standardization of surgical technique for appendectomy [6].

A uniform DPC would ideally be evidence-based. Unfortunately, limited evidence regarding outcomes specific to device utilization for laparoscopic appendectomy currently exists [7]. However, Miyano et al. have suggested a potential cost savings for use of pretied surgical loops rather than staples for controlling the base of the appendix [8]. There are no published reports describing differences in outcomes based upon the type of port, heat source or use of a specimen retrieval bag. Given this lack of evidence, we elected to design our uniform DPC to use low-cost ports, employ hook electrocautery in place of a pistol-grip heat source for division of the mesoappendix, and eliminate routine use of the specimen retrieval bag. We did not use reusable ports owing to institutional availability.

Table 3
Device utilization.

	Prior to uniform DPC, N = 346	With uniform DPC, N = 362	P value
	N (%)	N (%)	
Sheathed port(s)	344 (99.2%)	35 (9.6%)	<0.001
Low cost port(s)	0 (0.0%)	338 (92.6%)	<0.001
Stapler(s)	334 (94.4%)	74 (20.3%)	<0.001
Low cost loop(s)	0 (0.0%)	288 (79.7%)	<0.001
Pistol-grip heat source(s)	79 (22.3%)	14 (3.8%)	<0.001
Specimen retrieval bag	321 (90.7%)	220 (60.3%)	<0.001

Before implementation of the uniform DPC there was significant concern among the 6 pediatric surgeons that elimination of the specimen retrieval bag may increase the rate of surgical site infection and use of hook electrocautery to divide the mesoappendix would lead to bleeding complications. There were no returns to the operating room for bleeding in either group and no significant difference in the rate of surgical site infection was demonstrated.

There was a trend toward an increase in some of the measured complications including inpatient readmission, intraabdominal abscess, reoperation, and interventional radiology drainage with use of the uniform DPC. However, it is possible that these adverse events are representative of the learning curve that occurred as surgeons began using devices.

We anticipated a learning curve during the initial period of the transition to the unified DPC. There is a potential increase in the frequency of complications during the first 6 months after implementation. However, this learning curve did not manifest in a large increase in surgical times.

Limitations to this study include inadequate power to detect differences in complications such as intraabdominal abscess, subcutaneous abscess, reoperation, postoperative interventional radiology drainage, postoperative imaging, readmission, postdischarge ED evaluations.

Inclusion of patients with both preoperatively and postoperatively diagnosed urinary tract infections, pneumonia, and other potentially appendicitis-related issues dramatically increased the range of the secondary outcome data.

Reusable hook cautery use was not measured as there was no mechanism in place to document the use of devices found in our standard laparoscopy set.

We did not formally study operating room staff satisfaction with the uniform DPC.

4. Conclusion

We have demonstrated that implementation of a uniform DPC and technical standardization for laparoscopic appendectomy can significantly reduce cost. Furthermore, this can occur without dramatically increasing operative times, length of stay, or postoperative complications.

References

- [1] St Peter SD, Sharp SW, Holcomb III GW, et al. An evidence-based definition for perforated appendicitis derived from a prospective randomized trial. *J Pediatr Surg* 2008; 43:2242–5.
- [2] Standard cost manager: user's guide. Cleveland, Ohio: Intermountain Health Care Inc.; 1987.
- [3] Meier J, Duval M, Wilkes J, et al. Surgeon dependent variation in adenotonsillectomy costs in children. *Otolaryngol Head Neck Surg* 2014;150:887–92.
- [4] Drummond MF, Stoddart GL, Torrance GW. Methods for the economic evaluation of health care programmes. Oxford, United Kingdom: Oxford University Press; 1987.
- [5] Putnam LR, Levy SM, Johnson E, et al. Impact of a 24-hour discharge pathway on outcomes of pediatric appendectomy. *Surgery* 2014;156:455–61.
- [6] Avansino JR, Goldin AB, Risley R, et al. Standardization of operative equipment reduces cost. *J Pediatr Surg* 2013;48:1843–9.
- [7] Langer M, Safavi A, Skarsgard ED. Management of the base of the appendix in pediatric laparoscopic appendectomy: clip, ligate, or staple? *Surg Technol Int* 2013;23: 81–3.
- [8] Miyano G, Urao M, Lane GJ, et al. A prospective analysis of endoloops and endostaples for closing the stump of the appendix in children. *J Laparoendosc Adv Surg Tech A* 2011;21:177–9.