

Factors Influencing Aspiration During Swallowing in Healthy Older Adults

Susan G. Butler, PhD; Andrew Stuart, PhD; Xiaoyan Leng, PhD; Catherine Rees, MD;
Jeff Williamson, MD; Stephen B. Kritchevsky, PhD

Objectives/Hypothesis: Although flexible endoscopic evaluation of swallowing (FEES) is an established diagnostic tool, little data exist on the effects of varying liquid types on the swallowing outcomes in healthy older adults.

Study Design: Prospective.

Methods: Seventy-six healthy older adult volunteers participated (i.e., 18, 28, and 30 volunteers in the 7th, 8th, and 9th decades of life, respectively). The effects of age, sex, liquid type (i.e., water, skim, 2% milk, or whole milk), delivery method (i.e., cup or straw), and volume (i.e., 5, 10, 15, or 20 mL) on Penetration Aspiration Scale (PAS) scores were assessed during FEES.

Results: Penetration and silent aspiration were observed in 83% and 28% of the participants, respectively. This represented 19% and 3% of participants' 2,432 swallows, respectively. Liquid type ($P = .0001$), bolus volume ($P = .02$), and delivery method ($P = .04$) significantly affected PAS scores. PAS scores were significantly ($P < .05$) greater for milk versus water swallows, whole- versus skim-milk swallows, 10- and 20-mL versus 5-mL volumes, and straw versus cup delivery. The risk for aspiration increased by approximately two-, three-, and seven-fold with maximal increases in bolus volume, fat content of liquids, and age, respectively.

Conclusions: Occasional aspiration may be an underappreciated phenomenon during FEES in

healthy older adults. In addition, milk yielded a higher likelihood of aspiration than water. Accordingly, different liquid types, bolus volumes, and delivery methods should be used to ensure an accurate assessment of aspiration status in healthy older adults.

Key Words: Swallowing, aspiration, flexible endoscopic evaluation of swallowing, older adults, liquid, delivery method, bolus volume.

Level of Evidence: 2c

Laryngoscope, 120:2147–2152, 2010

INTRODUCTION

Flexible endoscopic evaluation of swallowing (FEES) is now routinely used to assess aspiration in outpatient and inpatient populations. Furthermore, the sensitivity of FEES is comparable, if not superior, to that of videofluoroscopic swallowing examination (VFSE) in the detection of aspiration.^{1–4} However, many aspects of the FEES protocol may vary among clinicians, and it is unclear how this variation might affect testing results. For example, bolus types are not standardized. Furthermore, we recently reported higher penetration aspiration scale (PAS) scores⁵ for milk versus water swallows in 14 healthy older adults [in review]. There was a significant difference between skim- and whole-milk PAS scores, suggesting that the higher the fat content of the liquid, the greater the risk for aspiration. If only water is tested during FEES, the risk for aspiration may be underappreciated. Likewise, if skim milk alone and not whole milk is tested, the risk for aspiration may also be underappreciated. In the same study [in review], we also found that bolus delivery (i.e., straw vs. cup delivery) and bolus size (i.e., the larger 20-mL bolus) yielded significantly more severe PAS scores.

Given the potential magnitude of our previous findings on FEES outcomes in healthy older adults, we sought to replicate and extend our previous pilot study with a larger sample size. The effects of age, sex, liquid type (i.e., water, skim milk, 2% milk, or whole milk), delivery method (i.e., cup or straw), and volume (i.e., 5, 10, 15, or 20 mL) on PAS scores were assessed during FEES. Thus, from previous research, we hypothesized

From the Department of Otolaryngology (S.G.B., C.R.); and Department of Biostatistical Sciences (X.L.), Department of Internal Medicine (J.W., S.B.K.), Wake Forest University School of Medicine, Winston-Salem, North Carolina, Department of Communication Sciences and Disorders, East Carolina University, Greenville, North Carolina (A.S.), U.S.A.

Editor's Note: This Manuscript was accepted for publication June 4, 2010.

Presented at the 18th Annual Meeting of the Dysphagia Research Society, San Diego, California, U.S.A., March 5–7, 2010.

This work was supported by NIDCD R03 DC009875, Wake Forest School of Medicine Claude D. Pepper Older Americans Independence Center (P30 AG21332), and the GCRC grant of Wake Forest University Baptist Medical Center (M01-RR07122). The authors have no other funding, financial relationships, or conflicts of interest to disclose.

Send correspondence to Susan G. Butler, PhD, CCC-LP, BRS-S, Associate Professor, Wake Forest University School of Medicine, Department of Otolaryngology, Medical Center Boulevard, Winston-Salem, NC 27157. E-mail: sbutler@wfubmc.edu

DOI: 10.1002/lary.21116

that liquids with higher fat content, larger volumes, and straw delivery would evoke higher PAS scores in healthy older adults.

MATERIALS AND METHODS

Participants

Seventy-six adults volunteered to participate in the study. They included 18 participants from 61 to 70 years old ($M = 65.9$ years, standard deviation [SD] = 1.9), 28 participants from 71 to 80 years old ($M = 77.6$ years, SD = 2.0), and 30 participants from 81 to 90 years old ($M = 83.8$ years, SD = 2.3). Participants reported no history of problems with swallowing, speech, or voice; and no known neurologic or otolaryngologic disorders. All participants were ambulatory and reported they were in good health. Participants were recruited by bulletins approved by the Wake Forest University Health Sciences Institutional Review Board. Informed consent was obtained.

Apparatus

A KayPENTAX Swallowing Workstation (KayPENTAX, Inc., Lincoln Park, NJ) with a PENTAX VNL 1070 STK nasopharyngo-laryngoscope and PENTAX EPK-1000 Digital Color Video Processors were used for the FEES.

Procedure

Participants underwent FEES while sitting in the upright position. A 3.1-mm digital flexible endoscope was lubricated with Surgilube (Altana Inc., Melville, NY) and passed transnasally, typically on the floor of the nose, by the first author to obtain a superior view of the hypopharynx. The endoscope was moved throughout the study between swallowing and postswallow positions to collect the data.

Swallowing position required that the distal end of the endoscope be just above the top of the epiglottis so that the entire base of the tongue, tip of the epiglottis, posterior pharyngeal wall, lateral pharyngeal walls (e.g., lateral channels), and laryngeal vestibule were visualized before bolus administration. The endoscope was maintained in swallowing position throughout bolus administration and was moved only while the scope was advanced to postswallow position following a bolus presentation. To obtain postswallow position, the distal end of the scope was advanced lower into the pharynx, past the tip of the epiglottis and into the upper portion of the laryngeal vestibule where the glottis and trachea below could be well visualized. Postswallow position was only held long enough to determine the PAS score, and then the scope was pulled back into swallowing position.

Four liquid boluses (i.e., water, skim milk, two percent milk, and whole milk) with four volumes (i.e., 5, 10, 15, and 20 mL) were administered with two delivery methods (i.e., straw vs. cup). All boluses were taken from the refrigerator simultaneously approximately 5 to 10 minutes before administration and were dyed with green food coloring to improve endoscopic visualization. Approximately 0.3 mL of green food coloring was added per 118 mL of liquid. The boluses were randomly presented to each participant in one data collection session of approximately 15 minutes. Premeasured liquid boluses were placed in 30-mL plastic cups for the cup delivery conditions. For straw administrations, a straw 185 mm in length with an inner diameter of 4.6 mm was placed in a 118-mL cone-shaped cup (Solo Cup Company, Urbana, IL). Participants were instructed before bolus administration that when they were handed a cup,

TABLE I.
Penetration Aspiration Scale Categorized According to Swallowing Status⁵.

Swallowing Status	PAS Score	Description
No penetration or aspiration		
	1	Material does not enter the airway.
Penetration	2	Material enters the airway, remains above the vocal folds, and is ejected from the airway.
	3	Material enters the airway, remains above the vocal folds, and is not ejected from the airway.
	4	Material enters the airway, contacts the vocal folds, and is ejected from the airway.
	5	Material enters the airway, contacts the vocal folds, and is not ejected from the airway.
Aspiration	6	Material enters the airway, passes below the vocal folds, and is ejected into the larynx or out of the airway.
	7	Material enters the airway, passes below the vocal folds, and is not ejected from the trachea despite effort.
	8	Material enters the airway, passes below the vocal folds, and no effort is made to eject.

PAS = penetration aspiration scale.

they should swallow all the liquid in one swallow when ready⁶; however, they could take more than one swallow if needed.

Swallows were reviewed in real time, slow motion, and frame by frame to assign the corresponding PAS score in accordance with previously published methods.^{5,7,8} Higher PAS scores reflect more abnormal swallows (Table I). A PAS score of 1 is a normal swallow with no material in the airway, scores between 2 and 5 indicate that material entered the laryngeal vestibule (i.e., penetration), and scores of 6 to 8 indicate that material passed below the vocal cords into the trachea (i.e., aspiration).

Data Analysis

A total of 2,432 data points were available for statistical analyses (i.e., 76 participants $\times$ 32 swallows). All swallows were scored with the eight-point PAS. A five-factor linear mixed model analysis of variance was used to examine the effect of liquid type, delivery type, bolus volume, sex, and age on the PAS scores. Since PAS scores were highly skewed, ranks were used in place of the PAS scores. A compound symmetry covariance structure was used to model the within-patient correlation. Linear contrasts were used to assess the pair-wise differences among the levels of significant main effects.

A logistic regression model was used to further investigate the relationship for penetration and aspiration status with predictor factors (i.e., liquid type, bolus volume, delivery method, and age). An odds ratio for each factor was obtained for penetration and aspiration, respectively, indicating the relative risk of penetration and aspiration between different factor levels. SAS version 9.1 (SAS Institute Inc., Cary, NC) was used for all

TABLE II.
Penetration Aspiration Scale Scores as a Function of Liquid Type, Delivery Method, and Bolus Volume.

Liquid	Delivery	Volume (mL)	PAS Score								M (SE)
			1	2	3	4	5	6	7	8	
Water	Straw	5	64	8	0	1	0	1	1	1	1.4 (0.1)
		10	66	7	1	0	1	0	1	0	1.3 (0.1)
		15	64	10	1	1	0	0	0	0	1.2 (0.1)
		20	63	11	1	0	0	0	1	0	1.3 (0.1)
	Cup	5	57	14	3	0	1	0	0	1	1.4 (0.1)
		10	64	10	0	0	1	0	0	1	1.3 (0.1)
		15	60	12	0	1	1	0	1	1	1.4 (0.1)
		20	62	12	0	1	0	0	0	1	1.3 (0.1)
Skim milk	Straw	5	61	13	1	0	0	0	1	0	1.3 (0.1)
		10	54	11	7	2	1	0	1	0	1.5 (0.1)
		15	55	12	5	1	1	0	2	0	1.5 (0.1)
		20	57	9	3	0	2	0	1	4	1.8 (0.2)
	Cup	5	68	5	2	0	0	0	0	1	1.2 (0.1)
		10	62	8	2	1	1	0	1	1	1.4 (0.1)
		15	61	9	2	1	0	0	1	2	1.5 (0.2)
		20	56	13	2	1	1	0	0	2	1.5 (0.1)
2% milk	Straw	5	57	10	3	0	5	0	0	1	1.6 (0.1)
		10	59	11	2	0	1	0	0	3	1.5 (0.2)
		15	54	13	1	0	1	0	4	2	1.8 (0.2)
		20	54	12	4	0	1	0	2	3	1.8 (0.2)
	Cup	5	61	6	3	1	4	0	0	1	1.5 (0.1)
		10	62	8	2	0	1	0	0	3	1.5 (0.2)
		15	57	9	3	0	1	1	0	5	1.8 (0.2)
		20	51	10	5	0	3	0	3	2	1.9 (0.2)
Whole milk	Straw	5	53	12	5	1	1	0	2	2	1.7 (0.2)
		10	55	15	4	0	0	0	1	1	1.5 (0.1)
		15	54	8	8	0	2	0	2	2	1.8 (0.2)
		20	53	9	7	0	0	1	1	5	1.9 (0.2)
	Cup	5	63	6	2	1	2	0	1	1	1.4 (0.1)
		10	62	7	5	1	0	0	0	1	1.4 (0.1)
		15	57	9	5	1	0	0	1	2	1.6 (0.2)
		20	56	8	7	1	1	0	1	2	1.6 (0.2)
Total			1,882	317	96	16	33	3	29	51	1.5 (0.03)

PAS = penetration aspiration scale; M = mean; SE = standard error.

analyses, and an α level of 0.05 was set for statistical significance.

RESULTS

PAS as Function of Liquid Type, Delivery Method, and Bolus Volume

PAS scores differed significantly by liquid type ($P = .0001$), bolus volume ($P = .02$), and delivery method ($P = .04$). In general, PAS scores were higher for milk swallows than water swallows, for larger versus smaller boluses, and for straw versus cup delivery. PAS scores for water ($M = 1.4$) differed significantly from those of skim milk ($M = 1.6$, $P = .03$), 2% milk ($M = 1.9$, $P = .0001$), and whole milk ($M = 1.8$, $P < .0001$). PAS scores for bolus volumes of 20 mL ($M = 1.8$) differed significantly from scores for 5 mL ($M = 1.5$, $P = .01$) and

10 mL ($M = 1.6$, $P = .01$). PAS scores as a function of liquid type, delivery method, and bolus volume collapsed across age and sex are shown in Table II.

Occurrences of and Sensorimotor Response to Penetration and Aspiration

There were 462 episodes (19%) of penetration and 83 episodes (3%) of aspiration during the 2,432 total swallows of the cohort. Eighty-three percent (63 of 76) and 28% (21 of 76) of the participants demonstrated penetration and silent aspiration, respectively, at some point in the testing. Eighty-five percent (123 of 145) and 61% (51 of 83) of penetration and aspiration episodes, respectively, did not elicit a sensorimotor response (i.e., throat clear and/or cough response). In other words, there was no attempt by the participant to clear the penetrated or

TABLE III.
Number of Aspiration Episodes as a Function of Number
of Participants.

Participants, No.	Silent Aspiration Episodes, No.
9	1
4	2
3	3
3	4
1	5
1	8

aspirated liquid. Of note, endoscopic evaluation scores transient penetration, which occurs with the swallow, as a PAS 2. That is, no penetration is seen in the laryngeal vestibule after the swallow, so no sensorimotor response would be expected. Accordingly, PAS 2 was eliminated from the sensorimotor calculations. The number of aspiration episodes as a function of the number of participants is presented in Table III. Observed PAS scores are summarized separately for each component of the testing condition in Table IV.

Odds Ratio of Penetration and Aspiration

Adjusted mean penetration and aspiration rates as a function of predictor factors (i.e., liquid type, bolus volume, delivery method, and age) are displayed with 95% confidence intervals in Table V. The risk of penetration increased with higher fat content of a liquid, larger bolus volume, and older age. Specifically, milk increased the risk for penetration by at least 30% as compared with water; the increase was 31% for skim milk ($P = .049$), 54% for 2% milk ($P = .007$), and 59% for whole milk ($P = .001$). A bolus volume of 20 mL increased the risk of penetration by 33% as compared with 5 mL ($P = .047$) and 10 mL ($P = .017$). Compared with 61 to 70

year olds, the 71 to 80 year olds had a doubled risk for penetration ($P = .044$), and the 81 to 90 year olds had a 2.5-fold increased risk ($P = .008$). Delivery type showed no statistical significance in regard to risk of penetration ($P = .062$). Similarly, higher fat content of a liquid and older age increased the risk of aspiration. Two-percent milk and whole milk increased the risk for aspiration by 3.2- and 2.7-fold, respectively, as compared with water. The aspiration risk increased by 6.7-fold in 81 to 90 year olds as compared with 61 to 70 year olds. Larger bolus volumes tended to increase the risk of aspiration ($P = .068$). Delivery type did not appear to have any effect on risk of aspiration ($P = .409$).

DISCUSSION

PAS scores as a function of liquid type (water, skim milk, 2% milk, whole milk), bolus volume (5, 10, 15, 20 mL), and bolus delivery method (straw, cup) were evaluated during FEES in 76 older healthy volunteers. Penetration and aspiration were observed in 19% and 3% of participant swallows, respectively. Twenty-eight percent (21 of 76) of the participants silently aspirated at some point in the testing, and 61% of the total aspiration occurrences did not elicit a cough reflex. PAS scores varied as a function of liquid type, with milk yielding more severe PAS scores than water. PAS scores also varied as a function of bolus volume and delivery method, with 20-mL bolus volumes and straw delivery yielding more severe PAS scores than small boluses and cup delivery. The risk of penetration increased with skim, 2%, and whole milk versus water; with 20-mL versus 5- and 10-mL boluses; and with 7th versus 8th and 9th decades of life. Similarly, the risk of aspiration increased with 2% and whole milk versus water, with skim versus 2% milk, and with 7th versus 9th decade of life.

These findings of higher PAS scores for milk versus water agree with previous findings from our pilot

TABLE IV.
Summary of Penetration Aspiration Scale Scores Across Testing Conditions.

Condition	No. of Swallows	M (SE)	Penetration (PAS 2–5)		Aspiration (PAS 6–8)		Silent Aspiration (PAS 8)	
			No. (%) of Swallows	% of Participants	No. (%) of Swallows	% of Participants	No. (%) of Swallows	% of Participants
Liquid								
Water	608	1.40 (0.05)	98 (16)	17.8	10 (2)	1.6	5 (1)	0.8
Skim milk	607	1.61 (0.07)	116 (19)	21.9	17 (3)	2.8	10 (2)	1.7
2% milk	605	1.88 (0.10)	120 (20)	24.8	26 (4)	5.0	20 (3)	3.3
Whole milk	607	1.81 (0.11)	128 (21)	25.4	28 (5)	4.3	16 (3)	2.6
Delivery								
Straw	1215	1.70 (0.08)	246 (20)	24.0	42 (3)	3.8	24 (2)	2.0
Cup	1212	1.65 (0.07)	216 (18)	20.9	37 (3)	3.1	27 (2)	2.0
Bolus volume, mL								
5	608	1.54 (0.07)	110 (18)	20.4	13 (2)	2.3	8 (1)	1.3
10	608	1.56 (0.09)	110 (18)	20.4	11 (2)	2.3	12 (2)	1.6
15	606	1.74 (0.08)	118 (19)	23.8	26 (4)	4.3	14 (2)	2.3
20	605	1.85 (0.09)	124 (20)	25.3	29 (5)	4.8	19 (3)	3.1

M = mean; SE = standard error; PAS = penetration aspiration scale.

TABLE V.
Adjusted Mean Penetration and Aspiration Rates From the Logistic Regression Model.

	Penetration			Aspiration		
	Rate, %	95% Confidence Limits		Rate, %	95% Confidence Limits	
Bolus volume, mL						
5	22.08	15.86	30.74	1.35	0.66	2.75
10	22.08	16	30.46	1.35	0.51	3.57
15	27	19.89	36.65	2.61	1.47	4.61
20	29.42	21.84	39.65	2.93	1.62	5.33
Delivery type						
Cup	22.74	17	30.41	1.72	0.98	3.01
Straw	27.37	20.48	36.57	2.17	1.06	4.46
Liquid type						
Water	18.64	13.06	26.6	0.98	0.48	1.99
Skim milk	24.38	17.91	33.19	1.7	0.83	3.51
2% milk	28.75	21.61	38.25	3.13	1.62	6.06
Whole milk	29.63	21.52	40.8	2.68	1.35	5.31
Age group, yr						
61–70	14.41	8.34	24.88	0.7	0.19	2.51
71–80	29.58	18.78	46.58	2.19	1.14	4.2
81–90	36.39	24.3	54.47	4.71	2.2	10.08

studies. Clinically, the choice to use water or milk during FEES may be an important one, especially in patient populations where the risk for aspiration pneumonia may be high (e.g., bedridden individuals). On the other hand, a few participants aspirated only water and not milk. Thus, these data suggest that testing both water and milk in clinical scenarios is prudent; however, if only one liquid type will be used for testing aspiration of liquids, milk will likely provide the highest yield of aspiration and risk for aspiration pneumonia.

All milk types elicited significantly higher PAS scores than water; however, the highest score was elicited by 2% milk; this finding may bear clinical significance. Logistic regression analyses yielded risks of aspiration that were approximately three times greater for 2% and whole milk compared to water and about two times greater for 2% versus skim milk. As we hypothesized, in general, the risk of penetration and aspiration increased with greater “miliness” of the liquid administered. Thus, it appears that if a clinician needs to rule out aspiration in an individual at high risk for aspiration pneumonia, testing with milk is prudent and likely testing with 2% milk is the best choice.

Testing with only milk during FEES could lead to an overly restrictive diet recommendation. Most participants who aspirated milk did not aspirate water. Thus, for patients with a low risk of pneumonia, our results suggest that it is important to test water as well as milk because the patient may not aspirate water, even if he or she demonstrates aspiration of milk.

Bolus size was a significant main effect in this study; 20 mL elicited higher PAS scores than 5 and 10 mL in healthy adults. In a previous report, an average bolus size was approximately 21 mL in individuals 70 to 79 years old⁹; thus, a 20-mL bolus is a reasonable

size for a participant to be expected to swallow. The risk of penetration also significantly increased with the 20-mL volume. Although all participants were instructed to swallow the entire amount in the cup in one swallow, several participants swallowed the 20-mL volumes in two swallows. Clinically, most professionals do not typically test with controlled bolus volumes; however, these findings show that if aspiration is seen on a larger bolus swallow, it might possibly be eliminated with a smaller one.

Mean PAS scores were found to be significantly higher for straw versus cup drinking. Although straw drinking elicited higher PAS scores, neither penetration nor aspiration were statistically more likely for straw versus cup drinking. So it appears that bolus delivery method marginally influenced measured swallowing outcomes in this sample of healthy adults. However, in a sample with dysphagia, the risk of penetration and aspiration may be significantly higher for straw versus cup drinking. Thus, bolus delivery methods warrant future systematic investigation in cohorts with dysphagia.

These data in healthy normal adults should be interpreted cautiously for patients who present clinically with aspiration risk and complication. For example, one could generalize these findings to patients with dysphagia. However, replication of this study in such patients would be needed to validate that generalization. Furthermore, although it is logical to associate more severe PAS scores with greater negative health consequences (e.g., pneumonia), there are no current data to demonstrate that direct association.

We found that participants in the 8th and 9th decades of life were twice as likely to demonstrate penetration as those in the 7th decade of life. Further,

individuals in their 9th decade of life were seven times more likely to aspirate than individuals in their 7th decade. This finding seems to reflect the likelihood that the aging process in the oropharynx and hypopharynx^{10,11} is more pronounced with increasing age.

The finding that 30% of this cohort silently aspirated replicates findings from the previous independent sample studies^{12,13} [in review]. But the participants who aspirated did not typically consistently aspirate, as only 3% of the study swallows yielded silent aspiration. A possible clinical interpretation of these findings is that an *isolated* event of liquid aspiration during a FEES procedure in a patient older than 70 years with dysphagia is within the range of normal and does not call for diet restrictions. However, clinical judgment should always be exercised relative to a patient's perceived risk of aspiration pneumonia and pulmonary health. Conversely, older adults who aspirate, even infrequently, may be at greater risk for community-acquired pneumonia or negative effects on the lungs. Even an isolated event of aspiration may provide a mechanism for direct transit of colonized pathogens (e.g., streptococci) from the oropharynx into the lungs.

Other researchers have demonstrated that FEES is more likely to identify penetration and aspiration than VFSE in patient populations,^{1,2,4,14} yet none have concurrently employed both techniques in the same sample of healthy adults. It is unknown whether the higher prevalence of aspiration in healthy older adults can be confirmed with VFSE.

A central question is whether all aspiration is bad in older adults. On the one hand, it may be important to identify healthy older adults who are at risk for aspiration, considering that its sequelae (e.g. community-acquired pneumonia, malnutrition, weight loss, and sarcopenia) are serious and can lead to functional decline and frailty. In that light, the most sensitive tool for detecting aspiration (e.g., FEES) should be employed whether evaluating healthy adults or patients. On the other hand, if aspiration is benign, either FEES or VFSE could be warranted, and greater sensitivity of one method versus another would be less pertinent. Currently, no data from longitudinal or cross-sectional studies exist to support or refute the association of aspiration with pneumonia in healthy older adults.

CONCLUSION

We found that varying bolus factors influence the outcomes of FEES. Milk elicited higher PAS scores and risk for penetration and aspiration as compared with water test boluses in a sample of healthy older adults. Bolus volumes of 20 mL yielded higher PAS scores and risk for penetration than those of 5 and 10 mL. Straw

delivery yielded higher PAS scores than cup delivery. Healthy participants in the 9th decade of life had a higher risk of penetration and aspiration than participants in the 7th decade of life. Our findings underscore the importance of choosing optimal test factors in regard to liquid type, bolus volume, and delivery methods during FEES; these choices could affect swallowing-safety diagnoses and diet recommendations.

Acknowledgment

We thank Karen Potvin Klein, MA, ELS (Research Support Core, Wake Forest University Health Sciences) for her editorial contributions to this manuscript.

BIBLIOGRAPHY

1. Rao N, Brady SL, Chaudhuri G, Donzelli JJ, Wesling MW. Gold-standard? Analysis of the videofluoroscopic and fiberoptic endoscopic swallowing examinations. *J Appl Res* 2003;3:89–96.
2. Leder SB, Sasaki CT, Burrell MI. Fiberoptic endoscopic evaluation of dysphagia to identify silent aspiration. *Dysphagia* 1998;13:19–21.
3. Wu CH, Hsiao TY, Chen JC, Chang YC, Lee SY. Evaluation of swallowing safety with fiberoptic endoscope: comparison with videofluoroscopic technique. *Laryngoscope* 1997; 107:396–401.
4. Langmore SE, Schatz K, Olson N. Endoscopic and videofluoroscopic evaluations of swallowing and aspiration. *Ann Otol Rhinol Laryngol* 1991;100:678–681.
5. Rosenbek JC, Robbins JA, Roecker EB, Coyle JL, Wood JL. A penetration-aspiration scale. *Dysphagia* 1996;11:93–98.
6. Daniels SK, Schroeder MF, DeGeorge PC, Corey DM, Rosenbek JC. Effects of verbal cue on bolus flow during swallowing. *Am J Speech Lang Pathol* 2007;16:140–147.
7. Colodny N. Interjudge and intrajudge reliabilities in fiberoptic endoscopic evaluation of swallowing (fees) using the penetration-aspiration scale: a replication study. *Dysphagia* 2002;17:308–315.
8. Daggett A, Logemann J, Rademaker A, Pauloski B. Laryngeal penetration during deglutition in normal subjects of various ages. *Dysphagia* 2006;21:270–274.
9. Adnerhill I, Ekberg O, Groher ME. Determining normal bolus size for thin liquids. *Dysphagia* 1989;4:1–3.
10. Nicosia MA, Hind JA, Roecker EB, et al. Age effects on the temporal evolution of isometric and swallowing pressure. *J Gerontol A Biol Sci Med Sci* 2000;55:M634–M640.
11. Crow HC, Ship JA. Tongue strength and endurance in different aged individuals. *J Gerontol A Biol Sci Med Sci* 1996;51:M247–M250.
12. Butler SG, Stuart A, Markley L, Rees C. Penetration and aspiration in healthy older adults as assessed during endoscopic evaluation of swallowing. *Ann Otol Rhinol Laryngol* 2009;118:190–198.
13. Butler SG, Stuart A, Kemp S. Flexible endoscopic evaluation of swallowing in healthy young and older adults. *Ann Otol Rhinol Laryngol* 2009;118:99–106.
14. Kelly AM, Drinnan MJ, Leslie P. Assessing penetration and aspiration: how do videofluoroscopy and fiberoptic endoscopic evaluation of swallowing compare? *Laryngoscope* 2007;117:1723–1727.